

On Minimal Length Factorizations of Finite Groups

María Isabel González Vasco, Martin Rötteler, and Rainer Steinwandt

CONTENTS

- 1. Introduction
- 2. Preliminaries
- 3. Some Examples and Construction Techniques
- 4. Simple Groups
- 5. Conclusions and Further Research
- Acknowledgments
- References

Logarithmic signatures are a special type of group factorizations, introduced as basic components of certain cryptographic keys. Thus, *short* logarithmic signatures are of special interest. We deal with the question of finding logarithmic signatures of *minimal length* in finite groups. In particular, such factorizations exist for solvable, symmetric, and alternating groups.

We show how to use the known examples to derive minimal length logarithmic signatures for other groups. Namely, we prove the existence of such factorizations for several classical groups and—in parts by direct computation—for all groups of order $< 175\,560$ ($= \text{ord}(J_1)$, where J_1 is Janko's first sporadic simple group). Whether there exists a minimal length logarithmic signature for each finite group still remains an open question.

1. INTRODUCTION

Public Key Cryptography nourishes on hard mathematical problems which very often, but not exclusively, arise from number theory. In the early '80s, several authors explored the possibility of using group theoretical problems for cryptography [Wagner and Magyarik 85, Wagner 90, Magliveras 86]. In particular owing to Magliveras et al., there are various proposals for cryptographic schemes which make use of special factorizations (so-called *logarithmic signatures*) of finite groups [Magliveras 86, Magliveras et al. 02]. Besides inspiring further cryptographic research [González Vasco and Steinwandt 02, González Vasco et al. 03, Birget et al. 02, Bohli et al. 02], these factorizations are interesting mathematical objects in themselves. For example, Hajós' work on Minkowski's conjecture illustrates that for abelian groups, this kind of factorization arises in the study of high-dimensional tilings (see [Stein and Szabó 94]).

Defined to be used as keys within a cryptographic scheme, the question of finding *short* logarithmic signatures arises naturally. The logarithmic signatures of abelian groups considered in Rédei's theorem (see, e.g., [Stein and Szabó 94]) are also examples of logarithmic signatures of minimal possible length.

2000 AMS Subject Classification: Primary 20D60; Secondary 20B05

Keywords: Logarithmic signatures, group factorizations, simple groups

In this contribution, we study the existence of *minimal length* logarithmic signatures for several families of finite groups: In Section 2, we give the basic definitions and notations; Section 3 details a (constructive) proof of the existence of minimal logarithmic signatures for solvable groups and recalls some other known constructions. Thereafter, we discuss how to extend these results to obtain new examples. From the arguments in Section 3, one can conclude that the smallest group for which a minimal length logarithmic signature does not exist (if there should be any!) must be a simple group. Hence, we devote Section 4 to studying several families of simple groups. We start by pointing out that there are suitable factorizations for $\mathrm{PSL}_2(q)$ and some other classical groups. Next we prove—in part through direct computation with a computer algebra system—that all finite groups of order $< 175\,560$ (the order of Janko’s first sporadic simple group) allow for a minimal length logarithmic signature. Whether such a factorization exists for all finite groups is to the best of our knowledge still an open problem. Some suggestions for possible directions for further research conclude the paper.

2. PRELIMINARIES

In a series of works, Magliveras et al. [Magliveras 02, Magliveras 86, Magliveras and Memon 92, Magliveras et al. 02] explored the possibility of building symmetric and asymmetric cryptosystems using certain group factorizations. We next recall the definition of logarithmic signatures, which are one of the mainstays of their research. For the basic related notions and results see, for instance, [Cusack 00].

Definition 2.1. Let G be a finite group. Next, denote by $\alpha = [\alpha_1, \dots, \alpha_s]$ a sequence of length $s \in \mathbb{N}_0$ such that each α_i ($1 \leq i \leq s$) is itself a sequence $\alpha_i = [\alpha_{i0}, \dots, \alpha_{ir_i-1}]$ with $\alpha_{ij} \in G$ ($0 \leq j < r_i$) and $r_i \in \mathbb{N}_0$. Then we call α a *logarithmic signature* for G if each $g \in G$ is represented uniquely as a product

$$g = \alpha_{1j_1} \cdots \alpha_{sj_s}$$

with $\alpha_{ij_i} \in \alpha_i$ ($1 \leq i \leq s$).

We refer to the sequences α_i , $i = 1, \dots, s$, as *blocks* of α and to the integer $\ell(\alpha) := \sum_{i=1}^s r_i$ as *length* of α .

Of course, for each group G there always exists a trivial logarithmic signature $\alpha := [[g \mid g \in G]]$ consisting of a single block. Being precise, we actually obtain $\mathrm{ord}(G)!$ “different” logarithmic signatures in this way, as a block

of a signature is an (ordered) sequence and not just a set. The distinction of whether a block is a sequence or a set is mainly motivated by the use of logarithmic signatures in the public key scheme MST_1 [Magliveras et al. 02], and as the subsequent discussion concerns only the length of logarithmic signatures, we can w.l.o.g. ignore this distinction.

Let us look at a more interesting example of a logarithmic signature: Assume we know a subgroup chain

$$G = G_0 > G_1 > \cdots > G_s = \{e_G\},$$

where e_G denotes the unit element in G . Now take $\alpha = [\alpha_i \mid i = 1, \dots, s]$, a sequence such that each $\alpha_i = [\alpha_{ij} \mid j = 0, \dots, [G_{i-1} : G_i] - 1]$ is a complete system of left coset representatives of G_{i-1} modulo G_i . It is easy to check that α is a logarithmic signature for G . Such logarithmic signatures are called *exact (left) transversal*. Analogously, one can use right coset representatives to derive *exact (right) transversal* logarithmic signatures.

In general, it is a nontrivial problem to check whether a group has logarithmic signatures of a fixed length. Luckily, we have at least a lower bound depending on the group order (first given in [González Vasco and Steinwandt 02]), which allows us to recognize the *shortest* ones:

Remark 2.2. Let G be a finite group and $\mathrm{ord}(G) = \prod_{j=1}^k p_j^{a_j}$ the prime factorization of the order of G (with $p_1, \dots, p_k$ different prime numbers). Moreover, denote by $\mathcal{B}(G) := \sum_{j=1}^k a_j p_j$ the sum of the prime divisors of $\mathrm{ord}(G)$ counting multiplicity. Then for each logarithmic signature α for G , we have

$$\ell(\alpha) \geq \mathcal{B}(G). \quad (2-1)$$

In the remaining part of the paper, we dwell on the problem of finding logarithmic signatures α for which the latter inequality is tight, i.e., where $\ell(\alpha) = \mathcal{B}(G)$ holds. Trivial examples are provided by the “empty” logarithmic signature $\alpha^{\{e\}} := []$ (of length 0) for the trivial group $\{e\}$ and by the logarithmic signature $\alpha^{C_p} := [[g \mid g \in C_p]]$ for a cyclic group C_p of prime order p . In the next section, we look at some more interesting examples.

3. SOME EXAMPLES AND CONSTRUCTION TECHNIQUES

We start by proving that the bound (2-1) is met for solvable groups (this result can also be found in [González Vasco and Steinwandt 02]).

Proposition 3.1. *Let G be a finite solvable group. Then there exists a logarithmic signature α for G of minimal length, i. e., such that $\ell(\alpha) = \mathcal{B}(G)$.*

Proof: Let G be a solvable group of order $\prod_{i=1}^k p_i$ with $p_1, \dots, p_k$ not necessarily distinct prime numbers. As G is solvable, it allows for a composition series

$$\gamma := G = G_0 > G_1 > \dots > G_k = \{e_G\}$$

where each $[G_{i-1} : G_i]$ is prime and $\prod_{i=1}^k [G_{i-1} : G_i] = \prod_{i=1}^k p_i$. Clearly, any exact transversal logarithmic signature derived from γ has length $\sum_{i=1}^k p_i$ and is thus of minimal length. $\square$

Remark 3.2. For abelian groups, there is an alternative construction which leads to a, in general, different, minimal length logarithmic signature: Let H be a cyclic group of order $\text{ord}(H) = p^m$ for some prime p , and take a any generator of H . Then $\alpha = [\alpha_1, \dots, \alpha_m]$ with $\alpha_i = [e_H, a^{p^{i-1}}, \dots, a^{(p-1)p^{i-1}}]$ ($i = 1, \dots, m$) is a logarithmic signature for H of length $m \cdot p = \mathcal{B}(H)$. Now just observe that the juxtaposition of minimal length logarithmic signatures for the cyclic p -primary factors of an abelian group G yields a minimal length logarithmic signature for G .

As neither the definition of logarithmic signatures nor the proof of the bound in Remark 2.2 exploits the fact that a group allows for inverse elements, it might be worth pointing out that for finite commutative monoids, minimal length factorizations do not always exist:

Example 3.3. Using a computer algebra system, one easily checks by exhaustive search that there exist no 3-element subsets α_1, α_2 of the monoid $(\mathbb{Z}/9\mathbb{Z}, \cdot, 1)$ such that $\alpha_1 \cdot \alpha_2 = \mathbb{Z}/9\mathbb{Z}$ holds.

Proofs of the tightness of the bound (2–1) for other families of groups often exploit peculiarities of the underlying group structure. This is the case for the symmetric and alternating groups; the existence of minimal length logarithmic signatures for the symmetric group S_n was first proven in [González Vasco and Steinwandt 02], and the tightness of the bound for the alternating group A_n was stated in [Magliveras 02]. Both proofs are constructive, and, moreover, in [Bohli et al. 02], there are examples of minimal length logarithmic signatures for S_n and A_n which are in addition tame and totally nontransversal (for the definitions of these notions, see [Magliveras et al. 02]).

The mentioned results for symmetric and alternating groups are in essence obtained by the same technique: Given a permutation representation of a group G , identify a point p so that its stabilizer G_p can be factored through a minimal length logarithmic signature and such that there exists a complete set of representatives of G modulo G_p which moves p cyclically. Actually, analogously as in Remark 3.2, the underlying idea is to factor the group into a “product of disjoint pieces” for which a minimal length logarithmic signature exists. For the case where the “disjoint pieces” are finite groups, we can formalize this idea through the concept of a *knit product* of groups [Michor 89] (also called *Zappa-Szép product*).

Definition 3.4. Let K and H be finite groups. Then we call two mappings $\kappa : H \times K \rightarrow K$, $\hbar : K \times H \rightarrow H$ an *automorphically knitted pair of actions* for (K, H) , provided that

1. the mapping

$$\begin{aligned} \psi_\kappa : H &\longrightarrow S_K \\ h &\longmapsto \kappa_h(\cdot) := \kappa(h, \cdot) \end{aligned}$$

is a group homomorphism,

2. the mapping

$$\begin{aligned} \phi_\hbar : K &\longrightarrow S_H \\ k &\longmapsto \hbar_k(\cdot) := \hbar(k, \cdot) \end{aligned}$$

is a group antihomomorphism, namely $\hbar_{e_K} = \text{id}_H$ and $\forall k_1, k_2 \in K : \hbar_{k_1} \hbar_{k_2} = \hbar_{k_2 k_1}$,

3. $\forall k_1, k_2 \in K, h \in H : \kappa_h(k_1 k_2) = \kappa_h(k_1) \kappa_{\hbar_{k_1}(h)}(k_2)$, and
4. $\forall k_1, k_2 \in K, h \in H : \hbar_k(h_1 h_2) = \hbar_{\kappa_{h_2}(k)}(h_1) \hbar_k(h_2)$,

where S_X denotes the group of permutations on the set X with functional composition as group operation, i. e., for $\sigma, \tau \in S_X$, we have $(\sigma\tau)(x) := \sigma(\tau(x))$.

Definition 3.5. Let K, H be groups, and $(\kappa, \hbar)$ an automorphically knitted pair of actions for (K, H) . Then the group defined over $K \times H$ with multiplication

$$(k_1, h_1) \cdot (k_2, h_2) := (k_1 \kappa_{h_1}(k_2), \hbar_{k_2}(h_1) h_2),$$

and unit element (e_K, e_H) is called *knit* or *Zappa-Szép product* of K and H and denoted by $K \times_{(\kappa, \hbar)} H$.

Note that $K \times \{e_H\}$ and $\{e_K\} \times H$ are subgroups of $K \times_{(\kappa, \hbar)} H$, isomorphic respectively to K and H . Also,

if $\psi_\kappa \equiv \text{id}_K$, then $\{e_K\} \times H$ is a normal subgroup of $K \times_{(\kappa, h)} H$, and thus we have a semidirect product (similarly, if $\phi_h \equiv \text{id}_H$). If both conditions hold, $K \times_{(\kappa, h)} H$ is a direct product of $K \times \{e_H\}$ and $\{e_K\} \times H$. Actually, the Zappa-Szép product generalizes the notions of direct and semidirect product, and it is easy to see that Zappa-Szép products can be used to derive groups that allow for a minimal length logarithmic signature:

Proposition 3.6. *Let G be a Zappa-Szép product of two finite groups K and H , and suppose there are logarithmic signatures α^K for K and α^H for H such that $l(\alpha^K) = \mathcal{B}(K)$ and $l(\alpha^H) = \mathcal{B}(H)$. Then there exists a logarithmic signature α^G for G with $l(\alpha^G) = \mathcal{B}(G)$.*

Proof: Let α^K and α^H be of the form

$$\alpha^K = [\alpha_1^K, \dots, \alpha_s^K], \alpha_i^K = [\alpha_{i0}^K, \dots, \alpha_{i r_i - 1}^K] \quad (i = 1, \dots, s),$$

$$\alpha^H = [\alpha_1^H, \dots, \alpha_t^H], \alpha_i^H = [\alpha_{i0}^H, \dots, \alpha_{i l_i - 1}^H] \quad (i = 1, \dots, t),$$

where $s, t \in \mathbb{N}_0$. Then clearly

$$\alpha^G := [\alpha_1^G, \dots, \alpha_s^G, \alpha_{s+1}^G, \dots, \alpha_{s+t}^G]$$

where

$$\begin{aligned} \alpha_i^G &:= [(\alpha_{i0}^K, e_H), \dots, (\alpha_{i r_i - 1}^K, e_H)] \quad (i = 1, \dots, s), \\ \alpha_{s+i}^G &:= [(e_K, \alpha_{i0}^H), \dots, (e_K, \alpha_{i l_i - 1}^H)] \quad (i = 1, \dots, t), \end{aligned}$$

is a minimal length logarithmic signature for G . $\square$

Therefore, the bound in Remark 2.2 is, in particular, tight for any group G that is an extension of two groups K and H which have logarithmic signatures of minimal length—and thus for all direct and semidirect products of groups with that property. The same applies to groups which are (set theoretically) decomposable as the product of two disjoint proper subgroups meeting the bound of Remark 2.2.

Example 3.7. In the following section, we see that the groups $\text{PSL}_2(q)$ have minimal length logarithmic signatures, and thus one concludes the tightness of (2–1) for the projective mock linear groups $\text{PML}_2(q)$ (where q is an even power of an odd prime), for $\text{PSL}_2(q)$ is a subgroup of $\text{PML}_2(q)$ of index two. For further discussion on projective mock linear groups, we refer to [Blackburn and Huppert 82, Chapter XI] and [Abhyankar 92].

Starting again with $\text{PSL}_2(q)$, one also verifies immediately the existence of minimal length logarithmic signatures for general linear groups $\text{GL}_2(q)$:

Example 3.8. As $\text{GL}_2(q)$ is a cyclic extension of $\text{SL}_2(q)$, and $\text{PSL}_2(q)$ is $\text{SL}(2, q)$ modulo its center, the bound (2–1) is tight for $\text{GL}_2(q)$, too.

Similarly, one easily sees that a finite group G has a minimal length logarithmic signature if it has a subnormal series whose factors do. Thus, for instance, the bound is tight for all nearly solvable groups.¹ As a result, if there exists any finite group which does *not* have a minimal length logarithmic signature, then the smallest (w.r.t. the group order) counterexample must be a simple group. In the remaining part of this paper, we therefore focus on simple groups.

4. SIMPLE GROUPS

Finite simple groups are indeed complex objects of study, but neatly classified [Gorenstein et al. 98] and relatively well understood. Actually, much research has been devoted to the problem of factoring finite simple groups as a product of two proper subgroups; for a detailed survey, see [Liebeck et al. 90]. However, the factorizations we are dealing with are significantly different: They are comprised of blocks which impose a unique factorization in the sense of Definition 2.1 on the group, but the blocks are not necessarily subgroups and in fact may have no “structure” at all. Of course, when trying to construct minimal length logarithmic signatures, we can, in particular, try to make use of known factorizations of finite simple groups and to exploit them for our purposes. To illustrate this approach, in the next section, we consider factorizations of simple groups into a product of Sylow subgroups.

4.1 Factoring into a Product of Sylow Subgroups

In [Holt and Rowley 93], Holt and Rowley study which groups G can be written as a product $P_1 \cdots P_k$ where $p_1, \dots, p_k$ are the different prime numbers dividing $\text{ord}(G)$ and each P_i is a p_i -Sylow subgroup of G . They show that such a factorization is possible for several types of groups, including $\text{PSL}_2(q)$ and $\text{PGL}_3(q)$ for any prime power q . As all Sylow subgroups are solvable and therefore meet the bound (2–1), we conclude the tightness of (2–1) for all simple groups $\text{PSL}_2(q)$ ($q > 3$) and for all simple groups $\text{PSL}_3(q)$ with $q \not\equiv 1 \pmod{3}$ (and hence, $\text{PSL}_3(q) \simeq \text{PGL}_3(q)$).²

¹A group is *nearly solvable* if it admits a subnormal series so that each of its factors is a Möbius group; all Möbius groups are solvable, except from A_5 .

²For $\text{PSL}_2(q)$, the existence of minimal length logarithmic signatures is also explored in [Magliveras 02]. However, the proof given

P_2	$= \langle (2, 15)(3, 10, 12, 7)(4, 11, 21, 13)(5, 17)(6, 8, 20, 9)(14, 18, 16, 19),$ $(2, 5)(3, 19, 8, 13)(4, 10, 14, 20)(6, 21, 7, 16)(9, 11, 12, 18)(15, 17),$ $(2, 5)(3, 8)(4, 16)(6, 20)(7, 10)(9, 12)(14, 21)(15, 17),$ $(3, 14)(4, 8)(6, 19)(7, 13)(9, 21)(10, 11)(12, 16)(18, 20) \rangle \leq \text{PSL}_3(4)$
P_3	$= \langle (1, 12, 5)(2, 8, 7)(3, 13, 15)(4, 10, 19)(6, 14, 18)(9, 16, 17),$ $(1, 5, 12)(2, 13, 9)(3, 17, 7)(6, 14, 18)(8, 15, 16)$ $(11, 21, 20) \rangle \leq \text{PSL}_3(4)$
P_5	$= \langle (1, 5, 17, 2, 15)(3, 21, 4, 8, 6)(7, 11, 12, 9, 19)$ $(13, 16, 20, 14, 18) \rangle \leq \text{PSL}_3(4)$
P_7	$= \langle (1, 20, 18, 10, 8, 12, 19)(2, 13, 6, 4, 17, 15, 3)$ $(5, 21, 16, 9, 14, 11, 7) \rangle \leq \text{PSL}_3(4)$

 TABLE 1. Factorization $\text{PSL}_3(4) = P_7 P_3 P_2 P_5$.

Also for $q \equiv 1 \pmod{3}$ such a factorization of $\text{PSL}_3(q)$ into a product of Sylow subgroups—and thereby a logarithmic signature of minimal possible length—can be available: Representing $\text{PSL}_3(4)$ as subgroup $\langle \alpha, \beta \rangle \leq S_{21}$ with generators

$$\begin{aligned} \alpha &= (4, 11, 20)(5, 15, 17)(6, 16, 18)(7, 14, 13) \\ &\quad (8, 12, 9)(10, 21, 19), \\ \beta &= (1, 8, 21, 16, 15, 3, 2)(4, 10, 20, 18, 17, 9, 7) \\ &\quad (5, 12, 11, 14, 19, 13, 6), \end{aligned}$$

and choosing Sylow subgroups P_2, P_3, P_5, P_7 as given in Table 1, we can express $\text{PSL}_3(4)$ as a product $\text{PSL}_3(4) = P_7 P_3 P_2 P_5$. At this, as always throughout the sequel, the product $(\sigma\pi) \in S_n$ of two permutations $\sigma, \pi \in S_n$ is understood to be the permutation that maps each $i \in \{1, \dots, n\}$ to $\pi(\sigma(i))$. The correctness of this factorization is easily checked by means of a computer algebra system like GAP [Team 97] or Magma [Bosma et al. 97].

Similar factorizations can be obtained for $\text{PSU}_4(2)$ and $\text{PSU}_3(4)$. Namely, representing $\text{PSU}_4(2)$ as subgroup $\langle \gamma, \delta \rangle \leq S_{45}$ with generators

$$\begin{aligned} \gamma &= (2, 5, 3)(4, 12, 7)(6, 17, 10)(8, 21, 14)(9, 19, 13) \\ &\quad (11, 29, 20)(16, 30, 25)(18, 28, 27)(22, 26, 32) \\ &\quad (23, 36, 31)(33, 35, 39)(34, 37, 41)(40, 42, 43), \\ \delta &= (1, 2, 4, 8, 15, 24)(3, 6, 11, 21, 31, 37) \\ &\quad (5, 9, 16, 14, 23, 34)(7, 13, 22, 33, 40, 20)(10, 18, 25) \\ &\quad (12, 17, 26, 35, 42, 30)(19, 28, 29)(32, 38, 43) \\ &\quad (36, 41, 45)(39, 44), \end{aligned}$$

and choosing Sylow subgroups P_2, P_3, P_5 as listed in Table 2, we can express $\text{PSU}_4(2)$ as a product $\text{PSU}_4(2) =$

there does not cover all choices of q . For example, for $\text{PSL}_2(13)$, no logarithmic signature can be obtained in this way.

$P_3 P_2 P_5$. Juxtaposing minimal length logarithmic signatures for these solvable factors yields the required minimal length logarithmic signature for $\text{PSU}_4(2)$.

Based on the representation $\text{PSU}_3(4) = \langle \epsilon, \zeta \rangle \leq S_{65}$ with

$$\begin{aligned} \epsilon &= (2, 9, 50, 12, 61, 38, 14, 3, 15, 4, 27, 63, 52, 23, 6) \\ &\quad (5, 32, 21, 10, 53, 33, 18, 11, 55, 45, 22, 47, 30, 16, 8) \\ &\quad (7, 62, 39, 29, 51, 25, 60, 17, 43, 42, 49, 44, 28, 34, 36) \\ &\quad (13, 48, 57, 59, 46, 41, 24, 20, 37, 54, 58, 35, 40, 19, 26) \\ &\quad (56, 64, 65), \\ \zeta &= (1, 2, 4, 8, 18, 39, 26, 48, 59, 44, 22, 21, 6, 15, 31) \\ &\quad (3, 7, 10, 23, 33, 47, 27, 16, 34, 9, 20, 41, 61, 40, 58) \\ &\quad (5, 12, 17, 37, 55, 54, 60, 38, 32, 56, 28, 52, 63, 62, 64) \\ &\quad (11, 25, 43, 45, 51, 46, 49, 50, 57, 65, 13, 14, 29, 19, 42) \\ &\quad (24, 35, 36), \end{aligned}$$

in Table 3, a respective factorization $\text{PSU}_3(4) = P_{13} P_5 P_2 P_3$ of $\text{PSU}_3(4)$ into Sylow subgroups is specified.

In general, we certainly cannot expect that a logarithmic signature of minimal length for a given finite (simple) group can be obtained by means of a factorization into Sylow subgroups. Actually, Holt and Rowley prove in [Holt and Rowley 93] that such a factorization does not exist for the simple group $\text{PSU}_3(3)$. At the moment, we do not know whether there exists a finite simple group—or equivalently any finite group—for which no logarithmic signature at all meeting the bound (2–1) exists. In the next section, we show that, in particular, for all sporadic simple Mathieu groups, a minimal length logarithmic signature exists, and thereafter, we prove that all simple groups of order $< 175\,560$ (the order of Janko’s first sporadic simple group) allow for a minimal length logarithmic signature, too. This implies that the bound

P_2	$\langle (1, 30, 45, 23)(2, 28, 3, 42)(4, 17, 16, 19)(5, 11)(6, 9, 26, 35)$ $(7, 43, 20, 27)(8, 31, 41, 12)(10, 44, 39, 18)(13, 40, 32, 38)$ $(14, 36, 37, 29)(15, 24, 34, 21)(22, 33),$ $(1, 15, 17, 6)(2, 44, 31, 13)(3, 32, 12, 18)(4, 40, 23, 39)(5, 22)(7, 20)$ $(8, 9, 42, 24)(10, 30, 38, 16)(11, 33)(14, 37)(19, 34, 45, 26)$ $(21, 28, 35, 41)(27, 29)(36, 43),$ $(1, 21, 17, 35)(2, 10, 31, 38)(3, 40, 12, 39)(4, 32, 23, 18)(5, 33)$ $(6, 41, 15, 28)(7, 20)(8, 26, 42, 34)(9, 19, 24, 45)(11, 22)$ $(13, 16, 44, 30)(14, 37)(27, 36)(29, 43) \rangle \leq \text{PSU}_4(2)$
P_3	$\langle (1, 27, 3, 19, 16, 12, 11, 34, 25)(2, 32, 26, 17, 8, 21, 33, 24, 36)$ $(4, 30, 22, 37, 43, 45, 10, 35, 31)(5, 41, 29)$ $(6, 20, 44, 28, 13, 14, 40, 38, 23)(7, 42, 18)(9, 39, 15),$ $(2, 44, 34)(3, 40, 24)(4, 22, 30)(5, 42, 41)(6, 32, 12)(7, 18, 39)$ $(8, 25, 28)(9, 15, 29)(10, 31, 35)(13, 20, 38)(14, 27, 17)(16, 33, 23)$ $(21, 36, 26)(37, 45, 43) \rangle \leq \text{PSU}_4(2)$
P_5	$\langle (1, 30, 40, 34, 20)(2, 42, 21, 18, 37)(3, 23, 10, 13, 7)(4, 35, 15, 27, 17)$ $(5, 11, 33, 22, 25)(6, 14, 45, 28, 9)(8, 38, 26, 36, 19)(12, 16, 24, 32, 43)$ $(29, 31, 41, 39, 44) \rangle \leq \text{PSU}_4(2)$

TABLE 2. Factorization $\text{PSU}_4(2) = P_3 P_2 P_5$.

(2–1) must be tight for arbitrary (not necessarily simple) groups of order $< 175\,560$.

4.2 Mathieu Groups

Among the sporadic simple groups, the five Mathieu groups M_{11} , M_{12} , M_{22} , M_{23} , and M_{24} were constructed two centuries ago. They are highly transitive permutation groups and can be obtained as the automorphism groups of Steiner systems. More precisely, M_{12} is the sharply 5-transitive automorphism group of a Steiner system $S(5, 6; 12)$ and M_{24} is the 5-transitive automorphism group of a Steiner system $S(5, 8; 24)$. We refer to [Beth et al. 99] for notations and the realization of the Mathieu groups as automorphism groups of Steiner systems. In the following, we employ a more direct construction which allows us to obtain the Mathieu groups as symmetries of the projective geometries $\text{PG}(2, \mathbb{F}_{11})$ and $\text{PG}(2, \mathbb{F}_{23})$, respectively.

We first outline an elementary construction of the Mathieu groups M_{11} and M_{12} based on the projective geometry $\text{PG}(2, \mathbb{F}_{11})$. Let

$$P = \{0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, \infty\}$$

denote the points of the standard coordinatization of $\text{PG}(2, \mathbb{F}_{11})$. Then M_{12} can be obtained as a permutation group on the set P . Namely, consider the following mappings (see [Gorenstein 82, Chapter 2.2]) which permute the set P :

$$f : x \mapsto x + 1, \quad g : x \mapsto -\frac{1}{x}, \quad h : x \mapsto 4x^2 - 3x^7.$$

Note that by convention $\infty = 1/0$ and $0 = 1/\infty$. The elements f and g generate the group $\text{PSL}_2(11)$ which is contained in $M_{12} := \langle f, g, h \rangle$. Explicitly, f , g , and h give rise to the following permutations σ , τ , and π of P :

$$\begin{aligned} \sigma &= (0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10), \\ \tau &= (0, \infty)(1, 10)(2, 5)(3, 7)(4, 8)(6, 9), \\ \pi &= (2, 6, 10, 7)(3, 9, 4, 5). \end{aligned}$$

Now M_{12} is a simple group of order 95 040 which is sharply 5-transitive on the points P . Furthermore, M_{11} is defined as the stabilizer of the point 0. Hence, $\text{ord}(M_{11}) = 7\,920$ and a direct calculation shows that

$$M_{12} = K_{12} \cdot M_{11}, \tag{4-1}$$

P_2	=	$\langle (1, 40, 29, 21)(2, 23, 15, 20)(3, 49, 34, 12)(4, 27, 46, 54)$ $(5, 50, 30, 26)(6, 32, 47, 38)(7, 35, 65, 25)(8, 41, 62, 13)$ $(9, 44, 36, 52)(10, 28, 45, 56)(11, 53, 59, 17)(14, 61, 24, 63)$ $(16, 19, 43, 37)(22, 60, 48, 31)(33, 39, 58, 55)(42, 51, 64, 57),$ $(1, 27, 24, 6)(2, 8, 22, 37)(3, 51, 52, 33)(4, 21, 38, 61)(5, 25, 45, 11)$ $(7, 50, 53, 56)(9, 64, 49, 39)(10, 59, 30, 35)(12, 55, 36, 42)$ $(13, 20, 43, 31)(14, 47, 29, 54)(15, 62, 48, 19)(16, 60, 41, 23)$ $(17, 28, 65, 26)(32, 63, 46, 40)(34, 57, 44, 58),$ $(1, 2, 14, 48)(3, 30, 44, 45)(4, 16, 32, 13)(5, 52, 10, 34)(6, 62, 54, 37)$ $(7, 42, 17, 39)(8, 27, 19, 47)(9, 28, 12, 50)(11, 58, 35, 51)$ $(15, 24, 22, 29)(20, 63, 60, 21)(23, 61, 31, 40)(25, 57, 59, 33)$ $(26, 36, 56, 49)(38, 41, 46, 43)(53, 55, 65, 64),$ $(1, 57, 14, 33)(2, 35, 48, 11)(3, 6, 44, 54)(4, 12, 32, 9)(5, 37, 10, 62)$ $(7, 31, 17, 23)(8, 30, 19, 45)(13, 26, 16, 56)(15, 25, 22, 59)$ $(20, 65, 60, 53)(21, 42, 63, 39)(24, 58, 29, 51)(27, 34, 47, 52)$ $(28, 41, 50, 43)(36, 46, 49, 38)(40, 64, 61, 55) \rangle \leq \text{PSU}_3(4)$
P_3	=	$\langle (1, 21, 22)(2, 14, 40)(3, 62, 42)(4, 65, 49)(5, 41, 25)(6, 33, 56)$ $(7, 9, 32)(8, 55, 44)(10, 13, 11)(12, 38, 53)(15, 29, 61)(16, 59, 30)$ $(17, 36, 46)(19, 39, 34)(23, 31, 60)(24, 63, 48)(26, 47, 51)$ $(27, 57, 28)(35, 45, 43)(37, 64, 52)(50, 54, 58) \rangle \leq \text{PSU}_3(4)$
P_5	=	$\langle (1, 40, 52, 42, 30)(2, 51, 60, 38, 8)(3, 55, 10, 24, 63)$ $(4, 37, 22, 33, 23)(5, 29, 21, 44, 64)(6, 53, 43, 28, 9)(7, 13, 26, 49, 27)$ $(12, 54, 65, 41, 50)(14, 61, 34, 39, 45)(15, 57, 31, 32, 62)$ $(16, 56, 36, 47, 17)(19, 48, 58, 20, 46),$ $(1, 47, 54, 5, 15)(2, 51, 60, 38, 8)(3, 53, 49, 34, 22)(4, 24, 9, 13, 14)$ $(6, 26, 61, 37, 63)(7, 45, 23, 10, 28)(11, 59, 25, 35, 18)$ $(12, 64, 62, 30, 36)(16, 41, 21, 31, 52)(17, 65, 29, 57, 40)$ $(19, 58, 46, 48, 20)(27, 39, 33, 55, 43)$ $(32, 42, 56, 50, 44) \rangle \leq \text{PSU}_3(4)$
P_{13}	=	$\langle (1, 3, 48, 30, 53, 6, 58, 18, 60, 56, 16, 55, 38)$ $(2, 65, 21, 61, 13, 46, 45, 19, 10, 59, 42, 8, 64)$ $(4, 54, 14, 50, 27, 44, 26, 15, 22, 34, 25, 29, 32)$ $(5, 47, 35, 28, 39, 51, 31, 37, 36, 9, 62, 33, 20)$ $(7, 49, 24, 43, 40, 57, 23, 63, 17, 52, 12, 41, 11) \rangle \leq \text{PSU}_3(4)$

TABLE 3. Factorization $\text{PSU}_3(4) = P_{13}P_5P_2P_3$.

where $K_{12} \leq M_{12}$ is the subgroup of order 12 generated by

$$\begin{aligned}\alpha &:= (0, 7)(1, 8)(2, 6)(3, \infty)(4, 10)(5, 9), \\ \beta &:= (0, 1)(2, 9)(3, 4)(5, 6)(7, 8)(10, \infty).\end{aligned}$$

In terms of the generators σ , τ , and π , we obtain the factorizations $\alpha = \pi\sigma^2\tau\pi^{-1}\tau\pi\tau^{-1}\sigma^{-2}\pi^{-1}$ and $\beta = \pi^{-1}\tau\sigma^2\tau^{-1}\sigma\pi$.

Since K_{12} is an abelian group isomorphic to a product $C_2 \times C_6$ of two cyclic groups, we obtain from Equation (4-1) that M_{12} has a logarithmic signature of minimal length if we can find one for M_{11} . To put it another way, we have that M_{12} is a Zappa-Szép product of K_{12} and M_{11} (see Section 3).

Focusing on M_{11} , we consider the stabilizer of the point 1 and obtain a group M_{10} of order 720 which is

known to have a composition series $\{e_{M_{10}}\} < A_6 < M_{10}$. Since A_6 allows for a logarithmic signature meeting the bound (2-1) [Magliveras 02, González Vasco and Steinwandt 02] and the factor group is isomorphic to C_2 , we obtain that M_{10} has a logarithmic signature of minimal length, and because of M_{11} being isomorphic to a Zappa-Szép product

$$M_{11} = C_{11} \cdot M_{10},$$

we can also construct minimal length logarithmic signatures for the small Mathieu groups M_{11} and M_{12} .

4.2.1 The Mathieu groups M_{22} , M_{23} , and M_{24} . Similarly to the construction of M_{12} , one may obtain M_{24} as a permutation group acting on the projective geometry $\text{PG}(2, \mathbb{F}_{23})$. This time, we start from $\text{PSL}_2(23)$ which

group	order	possible minimal length factorization
C_p (p prime)	p	trivial: $[[g \mid g \in C_p]]$
A_n ($n \geq 5$)	$n!/2$	stabilizer chain (see [Magliveras 02] and [Bohli et al. 02])
$\text{PSL}_2(q)$ ($q > 3$)	$\frac{q \cdot (q^2 - 1)}{\gcd(2, q - 1)}$	product of Sylow subgroups (Section 4.1)
$\text{PSL}_3(q)$ ($q \neq 3$)	$q^3(q^3 - 1)(q^2 - 1)$	product of Sylow subgroups (Section 4.1)
$\text{PSL}_3(4)$	20 160	product of Sylow subgroups (Table 1)
$\text{PSU}_4(2)$	25 920	product of Sylow subgroups (Table 2)
$\text{PSU}_3(4)$	62 400	product of Sylow subgroups (Table 3)
M_{11}	7 920	product of subgroups: $C_{11} \cdot A_6 \cdot C_2$
M_{12}	95 040	product of subgroups: $(C_2 \times C_6) \cdot M_{11}$
M_{22}	443 520	product of subgroups: $\text{PSL}_3(4) \cdot C_2 \cdot C_{11}$
M_{23}	10 200 960	product of subgroups: $C_{23} \cdot M_{22}$
M_{24}	244 823 040	product of subgroups: $S_4 \cdot M_{23}$

TABLE 4. Simple groups meeting the bound in Remark 2.2.

is generated by $f : x \mapsto x + 1$ and $g : x \mapsto -\frac{1}{x}$. The mapping, (see [Gorenstein 82])

$$h : x \mapsto -3x^{15} + 4x^4,$$

defines a permutation of the points of $\text{PG}(2, \mathbb{F}_{23})$. The Mathieu group M_{24} is defined as $M_{24} := \langle \sigma, \tau, \pi \rangle$, where the permutations σ, τ , and π are obtained from f, g , and h . Explicitly, we have

$$\begin{aligned} \sigma &= (0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, \\ &\quad 16, 17, 18, 19, 20, 21, 22), \\ \tau &= (0, \infty)(1, 22)(2, 11)(3, 15)(4, 17)(5, 9)(6, 19) \\ &\quad (7, 13)(8, 20)(10, 16)(12, 21)(14, 18), \\ \pi &= (2, 16, 9, 6, 8)(3, 12, 13, 18, 4)(7, 17, 10, 11, 22) \\ &\quad (14, 19, 21, 20, 15). \end{aligned}$$

Note that M_{24} can be written as a Zappa-Szép product in the following way:

$$M_{24} = S_4 \cdot M_{23},$$

where M_{23} denotes the stabilizer of the point 0 in M_{24} . The symmetric group $S_4 := \langle \alpha, \beta \rangle \leq M_{24}$ is generated by the permutations

$$\begin{aligned} \alpha &:= (0, 2)(1, 14)(3, 8)(4, \infty)(5, 20)(6, 17)(7, 10) \\ &\quad (9, 21)(11, 18)(12, 16)(13, 15)(19, 22), \\ \beta &:= (0, 1, \infty, 12)(2, 7, 3, 9)(4, 18, 13, 22)(5, 10, 16, 19) \\ &\quad (6, 11, 14, 21)(8, 20, 15, 17). \end{aligned}$$

We can descend one more step to M_{22} since the corresponding transversal can be chosen to be a cyclic group of order 23. More precisely, we have $M_{23} = C_{23} \cdot M_{22}$, where M_{22} stabilizes the points 0 and 1, and C_{23} is a

cyclic group of order 23. In fact, as generator of C_{23} , we can choose an arbitrary element of order 23 in M_{23} .

Finally, M_{22} can be realized as follows: We define the subgroups

$$\begin{aligned} C_{11} &:= \langle (2, 19, 9, 7, 18, 12, \infty, 5, 11, 16, 21) \\ &\quad (3, 17, 8, 10, 13, 20, 22, 14, 6, 4, 15) \rangle \leq M_{22} \end{aligned}$$

and

$$\begin{aligned} C_2 &:= \langle (2, 10)(3, 19)(4, 11)(7, 13)(8, 21)(9, 15)(12, 14) \\ &\quad (16, 17) \rangle \leq M_{22}. \end{aligned}$$

Then the subset $A := C_2 \cdot C_{11}$ of M_{22} has the property that for each point $x \in \{2, \dots, 22, \infty\}$, there is an element $\rho \in A$ such that $\rho(2) = x$. On the other hand, the stabilizer of the points 0, 1, and 2 inside M_{24} is isomorphic to the simple group $\text{PSL}_3(4)$ of order 20 160. Thus, using the result that $\text{PSL}_3(4)$ has a logarithmic signature of minimal possible length (see Table 1), we also obtain a logarithmic signature for $M_{22} = \text{PSL}_3(4) \cdot A$ that meets the bound (2–1). Hence, by the previous arguments, M_{23} and M_{24} also have logarithmic signatures of minimal possible length.

4.3 A Lower Bound for the Size of a Counterexample

In Table 4, the simple groups for which we already know that the bound in Remark 2.2 is tight are listed.

In particular, this list covers all simple groups of order $\leq 2^{17}$ up to the following three exceptions:

- $\text{PSU}_3(3)$ (of order 6 048)

C_4	$= \langle (1, 7, 13, 21)(2, 8, 19, 4)(3, 24, 22, 18)(5, 6, 10, 14)(9, 23)(11, 20, 28, 15)(12, 16)(17, 27, 25, 26) \rangle \leq \text{PSU}_3(3)$
C_7	$= \langle (1, 11, 8, 14, 20, 27, 2)(3, 5, 12, 21, 18, 4, 28)(6, 23, 24, 26, 22, 9, 13)(7, 25, 16, 17, 10, 15, 19) \rangle \leq \text{PSU}_3(3).$

TABLE 5. Factors C_4 and C_7 of the factorization $\text{PSU}_3(3) = G_1 C_4 C_7$.

C_5	$= \langle (1, 38, 43, 40, 64)(2, 50, 32, 63, 44)(3, 18, 10, 65, 61)(4, 7, 15, 24, 12)(5, 19, 33, 54, 17)(6, 8, 36, 27, 13)(9, 31, 48, 47, 30)(11, 49, 51, 20, 23)(14, 60, 16, 53, 57)(21, 39, 56, 35, 37)(22, 46, 58, 45, 59)(25, 34, 62, 26, 41)(28, 55, 29, 42, 52) \rangle \leq \text{Sz}(8)$
C_{13}	$= \langle (1, 46, 21, 18, 47, 51, 22, 31, 15, 4, 39, 6, 41)(2, 56, 32, 27, 5, 49, 3, 20, 40, 54, 50, 59, 23)(7, 42, 48, 61, 37, 65, 53, 36, 19, 8, 17, 45, 43)(9, 52, 55, 29, 38, 12, 62, 25, 57, 24, 11, 58, 30)(10, 63, 26, 14, 44, 35, 13, 34, 33, 16, 64, 28, 60) \rangle \leq \text{Sz}(8)$

TABLE 6. Factors C_5 and C_{13} of the factorization $\text{Sz}(8) = G_1 C_5 C_{13}$.

- $\text{Sz}(8)$ (of order 29 120)

- $\text{PSU}_3(5)$ (of order 126 000)

Subsequently, we give a logarithmic signature for each of these three groups. They have been found and verified by means of the computer algebra system Magma.

4.3.1 $\text{PSU}_3(3)$. We represent $\text{PSU}_3(3)$ as a subgroup $\langle \alpha, \beta \rangle \leq S_{28}$ with generators

$$\begin{aligned} \alpha &= (2, 7, 23, 26, 17, 13, 6, 3)(4, 19, 11, 28, 25, 24, 16, 9) \\ &\quad (5, 18, 10, 14, 15, 8, 20, 12)(21, 27), \\ \beta &= (1, 2, 4, 10, 8, 16, 13, 22)(3, 7, 9, 17, 6, 12, 21, 5) \\ &\quad (11, 19, 26, 14, 15, 23, 24, 25)(27, 28). \end{aligned}$$

Then $\text{PSU}_3(3)$ can be factored into a product $\langle \alpha, \beta \rangle = G_1 C_4 C_7$ where G_1 is the stabilizer of 1, and C_4 , respectively, C_7 , is a cyclic group of order 4, respectively, 7. Precise choices for C_4 and C_7 are given in Table 5.

As the stabilizer G_1 (of order $216 = \text{ord}(\text{PSU}_3(3))/(4 \cdot 7)$) is solvable, we immediately obtain a minimal length logarithmic signature for $\text{PSU}_3(3)$ by juxtaposing minimal length logarithmic signatures for the three subgroups G_1 , C_4 , and C_7 .

4.3.2 $\text{Sz}(8)$. We represent the Suzuki group $\text{Sz}(8)$ as subgroup $\langle \alpha, \beta \rangle \leq S_{65}$ with generators

$$\begin{aligned} \alpha &= (1, 2)(3, 4)(5, 7)(6, 9)(8, 12)(10, 13)(11, 15)(14, 19) \\ &\quad (16, 21)(17, 23)(18, 25)(20, 28)(22, 31)(24, 33) \\ &\quad (26, 35)(27, 32)(29, 37)(30, 39)(34, 43)(36, 46) \\ &\quad (38, 48)(41, 51)(42, 44)(45, 55)(47, 50)(49, 58) \\ &\quad (52, 60)(53, 61)(54, 59)(56, 62)(57, 63)(64, 65), \\ \beta &= (1, 3, 5, 8)(4, 6, 10, 14)(7, 11, 16, 22)(9, 12, 17, 24) \\ &\quad (13, 18, 26, 36)(15, 20, 29, 38)(19, 27, 31, 28) \\ &\quad (21, 30, 40, 50)(23, 32, 41, 52)(25, 34, 44, 54) \\ &\quad (33, 42, 53, 43)(35, 45, 56, 63)(37, 47, 51, 46) \\ &\quad (39, 49, 59, 60)(48, 57, 55, 58)(61, 64, 62, 65). \end{aligned}$$

Then $\text{Sz}(8)$ can be factored into a product $\text{Sz}(8) = G_1 C_5 C_{13}$ where G_1 is the stabilizer of 1, and C_5 , respectively, C_{13} , is cyclic of order 5, respectively, 13. Concrete choices for the cyclic groups C_5 and C_{13} are given in Table 6.

As the stabilizer G_1 (of order $448 = \text{ord}(\text{Sz}(8))/(5 \cdot 13)$) is solvable, we immediately obtain the required minimal length logarithmic signature for $\text{Sz}(8)$ by juxtaposing minimal length logarithmic signatures for the three factors G_1 , C_5 , and C_{13} .

C_2	$= \langle (1, 60)(3, 6)(4, 27)(5, 21)(7, 94)(8, 59)(9, 12)(10, 83)(11, 42)(13, 30)(14, 84)(15, 58)(16, 47)(17, 32)(18, 74)(19, 115)(20, 77)(22, 54)(23, 78)(24, 120)(25, 82)(26, 43)(28, 114)(29, 100)(31, 76)(33, 106)(34, 35)(36, 45)(37, 80)(38, 39)(40, 105)(41, 98)(44, 55)(46, 67)(48, 79)(49, 116)(50, 61)(52, 124)(53, 71)(56, 91)(57, 70)(62, 97)(63, 119)(64, 92)(65, 88)(66, 111)(68, 103)(69, 109)(72, 96)(73, 101)(75, 126)(81, 122)(86, 107)(87, 113)(90, 110)(93, 108)(95, 125)(102, 121)(104, 112)(118, 123) \rangle \leq \text{PSU}_3(5)$
$C_3^{(1)}$	$= \langle (1, 81, 13)(2, 119, 26)(3, 16, 105)(4, 121, 97)(5, 120, 109)(6, 24, 113)(7, 10, 28)(8, 40, 74)(9, 66, 94)(11, 39, 49)(12, 112, 62)(14, 60, 83)(15, 126, 122)(17, 123, 56)(18, 76, 33)(19, 69, 91)(20, 101, 45)(21, 87, 61)(22, 78, 104)(23, 100, 65)(25, 43, 72)(27, 67, 75)(29, 58, 36)(30, 51, 38)(31, 118, 108)(32, 59, 89)(34, 41, 55)(35, 82, 71)(37, 103, 110)(42, 63, 86)(44, 114, 107)(46, 48, 116)(47, 85, 115)(50, 90, 80)(52, 77, 70)(53, 124, 111)(54, 98, 102)(57, 84, 117)(64, 79, 88)(68, 93, 106)(73, 95, 99)(92, 125, 96) \rangle \leq \text{PSU}_3(5)$
$C_3^{(2)}$	$= \langle (1, 8, 6)(2, 57, 56)(3, 88, 104)(4, 79, 9)(5, 22, 97)(7, 30, 67)(10, 58, 54)(11, 28, 27)(12, 75, 106)(13, 19, 94)(14, 50, 51)(15, 68, 66)(16, 34, 44)(17, 36, 73)(18, 25, 32)(20, 93, 112)(21, 26, 70)(23, 87, 24)(29, 64, 111)(31, 47, 116)(33, 105, 91)(35, 98, 84)(37, 72, 121)(38, 83, 40)(39, 59, 78)(41, 110, 71)(42, 43, 69)(45, 108, 123)(46, 95, 102)(48, 100, 77)(49, 115, 55)(52, 80, 63)(53, 125, 96)(60, 82, 89)(61, 124, 76)(62, 113, 74)(65, 126, 117)(81, 114, 118)(85, 90, 119)(86, 120, 92)(99, 109, 101)(103, 107, 122) \rangle \leq \text{PSU}_3(5)$
C_7	$= \langle (1, 108, 125, 9, 17, 95, 64)(2, 39, 63, 34, 111, 25, 60)(3, 33, 68, 30, 74, 47, 107)(4, 5, 32, 52, 67, 57, 62)(6, 23, 78, 28, 80, 46, 122)(7, 24, 50, 100, 48, 104, 15)(8, 18, 85, 93, 44, 51, 16)(10, 116, 113, 22, 90, 126, 65)(11, 55, 72, 26, 42, 124, 14)(12, 77, 121, 75, 120, 84, 59)(13, 92, 56, 88, 118, 94, 73)(19, 36, 45, 103, 35, 21, 54)(20, 98, 110, 69, 82, 29, 87)(27, 112, 109, 70, 117, 97, 89)(31, 99, 96, 79, 66, 81, 123)(37, 61, 91, 101, 71, 102, 58)(38, 115, 105, 106, 40, 114, 76)(41, 119, 53, 49, 43, 86, 83) \rangle$

TABLE 7. Cyclic factors of the factorization $\text{PSU}_3(5) = C_7 C_3^{(1)} C_3^{(2)} C_2 G_1$.

4.3.3 $\text{PSU}_3(5)$. We represent $\text{PSU}_3(5)$ as a subgroup $\langle \alpha, \beta \rangle \leq S_{126}$ with generators

$$\begin{aligned} \alpha = & (1, 2, 4, 10, 25)(3, 7, 17, 41, 19)(5, 13, 33, 68, 111) \\ & (6, 14, 36, 34, 70)(8, 20, 47, 81, 48)(9, 22, 32, 61, 106) \\ & (11, 28, 37, 40, 82)(12, 30, 24, 55, 65) \\ & (15, 38, 77, 29, 64)(16, 39, 43, 85, 73) \\ & (18, 44, 75, 118, 109)(21, 49, 93, 126, 120) \\ & (23, 53, 42, 83, 52)(26, 58, 104, 122, 101) \\ & (27, 60, 105, 110, 124)(31, 35, 72, 116, 66) \\ & (45, 87, 123, 108, 96)(46, 89, 95, 100, 59) \\ & (50, 94, 57, 76, 103)(51, 97, 114, 71, 107) \end{aligned}$$

$$\begin{aligned} & (54, 88, 74, 63, 98)(56, 102, 117, 99, 121) \\ & (62, 86, 78, 90, 80)(67, 79, 112, 92, 125) \\ & (69, 113, 84, 115, 119), \\ \beta = & (2, 13, 112, 108, 24, 93, 45, 7) \\ & (3, 32, 53, 102, 54, 65, 75, 16) \\ & (4, 41, 52, 104, 11, 68, 99, 30) \\ & (5, 81, 122, 18, 56, 34, 19, 63) \\ & (6, 17, 72, 69, 101, 10, 55, 31) \\ & (8, 29, 100, 40, 84, 67, 37, 57) \\ & (9, 36, 98, 124, 27, 94, 117, 66) \\ & (12, 120, 42, 43, 33, 15, 26, 106) \end{aligned}$$

(14, 76, 64, 85, 86, 96, 70, 97)
 (20, 73, 82, 83, 125, 50, 39, 113)
 (21, 74, 87, 23, 59, 105, 110, 61)
 (22, 60, 88, 123, 78, 62, 90, 116)
 (28, 118, 77, 79, 44, 46, 115, 121)
 (35, 80, 109, 48, 95, 49, 114, 126)
 (38, 91, 47, 92, 51, 107, 71, 119)
 (58, 103, 89, 111).

Then $\text{PSU}_3(5)$ can be factored into a product $\langle \alpha, \beta \rangle = C_7 C_3^{(1)} C_3^{(2)} C_2 G_1$ where G_1 is the stabilizer of 1, C_2 is a cyclic group of order 2, both $C_3^{(1)}$ and $C_3^{(2)}$ are cyclic groups of order 3, and C_7 is a cyclic group of order 7. Precise choices for the four cyclic factors are given in Table 7. As the stabilizer G_1 (of order $1000 = \text{ord}(\text{PSU}_3(5))/(2 \cdot 3 \cdot 3 \cdot 7)$) is solvable, we immediately obtain a minimal length logarithmic signature for $\text{PSU}_3(5)$ by juxtaposing minimal length logarithmic signatures for the five subgroups G_1 , C_2 , $C_3^{(1)}$, $C_3^{(2)}$, and C_7 .

Thus, if there is any finite group G such that no logarithmic signature for G meets the bound in Remark 2.2, then the smallest counterexample (w.r.t. the group order) has to be both simple and of order $> 2^{17}$. Further on, we can exclude all simple groups covered by Table 4, and thus the smallest group for which the tightness of (2-1) is open is Janko's first sporadic group J_1 of order 175 560.

5. CONCLUSIONS AND FURTHER RESEARCH

Motivated by the question of finding short keys for the public key cryptosystem MST_1 , we have shown that various finite groups allow for logarithmic signatures of minimal possible length. Unfortunately, so far we could not answer the question whether there is any finite group for which the bound (2-1) is not tight, but we have shown that there can be no such group of cardinality smaller than Janko's first sporadic simple group.

Both from the mathematical and the cryptographic point of view, it would be desirable to identify further families of—not necessarily simple—groups for which the bound (2-1) is tight. On the cryptographic side, the question also arises whether logarithmic signatures of minimal length can be found, where effectively factoring group elements along the logarithmic signature is computationally hard. Such logarithmic signatures would be desirable for realizing the MST_1 public key cryptosystem.

ACKNOWLEDGEMENTS

We thank Consuelo Martínez for helpful comments and discussions. The work of the first author was partially supported by project BFM2001-3239-C03-01.

REFERENCES

- [Abhyankar 92] S. S. Abhyankar. “Galois Theory on the Line in Nonzero Characteristic.” *Bulletin of the American Mathematical Society* 27:1 (1992), 68–133.
- [Beth et al. 99] Th. Beth, D. Jungnickel, and H. Lenz. *Design Theory*, Volume I, Second edition. Cambridge, UK: Cambridge University Press, 1999.
- [Birget et al. 02] J.-C. Birget, S. S. Magliveras, and W. Wei. “Trap Doors from Subgroup Chains and Recombinant Bilateral Transversals. In *Actas de la VII Reunión Española de Criptología y Seguridad de la Información; Tomo I*, edited by S. González Jiménez and C. Martínez López, pp. 31–48, Oviedo: Servicio de Publicaciones, Universidad de Oviedo, 2002.
- [Blackburn and Huppert 82] N. Blackburn and B. Huppert. *Finite Groups III. Die Grundlehren der Mathematischen Wissenschaften*. Berlin-New York: Springer-Verlag, 1982.
- [Bohli et al. 02] J.-M. Bohli, M. I. González Vasco, C. Martínez, and R. Steinwandt. “Weak Keys in MST_1 .” *Cryptology ePrint Archive*, Report 2002/070, 2002.
- [Bosma et al. 97] W. Bosma, J. Cannon, and C. Playoust. “The Magma Algebra System I: The User Language.” *Journal of Symbolic Computation* 24 (1997), 235–265.
- [Cusack 00] C. A. Cusack. “Group Factorizations in Cryptography.” PhD thesis, University of Nebraska, 2000.
- [González Vasco et al. 03] M.I. González Vasco, C. Martínez, and R. Steinwandt. “Towards a Uniform Description of Several Group Based Cryptographic Primitives.” To appear in *Designs, Codes and Cryptography*, 2003.
- [González Vasco and Steinwandt 02] M. I. González Vasco and R. Steinwandt. “Obstacles in Two Public-Key Cryptosystems Based on Group Factorizations.” In *Cryptology*, edited by K. Nemoga and O. Grošek, pp. 23–37, Tatra Mountains Mathematical Publications, Volume 25. Bratislava: Mathematical Institute, Slovak Academy of Sciences, 2002.
- [Gorenstein 82] D. Gorenstein. *Finite Simple Groups*. University Series in Mathematics. New York: Plenum Press, 1982.
- [Gorenstein et al. 98] D. Gorenstein, R. Lyons, and R. Solomon. *The Classification of the Finite Simple Groups*, Mathematical Surveys and Monographs, Volume 40(1). Providence, RI: AMS 1998.
- [Holt and Rowley 93] D. F. Holt and P. Rowley. “On Products of Sylow Subgroups in Finite Groups.” *Archiv der Mathematik* 60:2 (1993), 105–107.

- [Liebeck et al. 90] M. W. Liebeck, C. E. Praeger, and J. Saxl. *The Maximal Factorizations of the Finite Simple Groups and their Automorphism Groups*, Memoirs of the AMS, Volume 86(432). Providence, RI: AMS, 1990.
- [Magliveras 86] S. S. Magliveras. “A Cryptosystem from Logarithmic Signatures of Finite Groups.” In *Proceedings of the 29th Midwest Symposium on Circuits and Systems*, pp. 972–975. Amsterdam: Elsevier Publishing Company, 1986.
- [Magliveras 02] S. S. Magliveras. “Secret- and Public-key Cryptosystems from Group Factorizations.” In *Cryptography*, edited by K. Nemoga and O. Grošek, pp. 11–22, Tatra Mountains Mathematical Publications, Volume 25. Bratislava: Mathematical Institute, Slovak Academy of Sciences, 2002.
- [Magliveras and Memon 92] S. S. Magliveras and N. D. Memon. “Algebraic Properties of Cryptosystem PGM.” *Journal of Cryptology* 5 (1992), 167–183.
- [Magliveras et al. 02] S. S. Magliveras, D. R. Stinson, and T. van Trung. “New Approaches to Designing Public Key Cryptosystems Using One-Way Functions and Trapdoors in Finite Groups.” *Journal of Cryptology* 15:4 (2002), 285–297.
- [Michor 89] P. W. Michor. “Knit Products of Graded Lie Algebras and Groups.” In *Proceedings of the Winter School on Geometry and Physics, Srni 1988, Ser. II, 22*, pp. 171–175, Palermo: Suppl. Rendiconti Circolo Matematico di Palermo, 1989.
- [Stein and Szabó 94] S. K. Stein and S. Szabó. *Algebra and Tiling. Homomorphisms in the Service of Geometry*. The Carus Mathematical Monographs, No. 25. Washington, DC: The Mathematical Association of America, 1994.
- [Team 97] The GAP Team. “GAP—Groups, Algorithms, and Programming.” Lehrstuhl D für Mathematik, RWTH Aachen, Germany and School of Mathematical and Computational Sciences, Univ. St. Andrews, Scotland, 1997.
- [Wagner 90] N. R. Wagner. “Searching for Public-Key Cryptosystems.” In *Proceedings of the 1984 Symposium on Security and Privacy (SSP '84)*, pp. 91–98. Los Alamitos, CA: IEEE Computer Society Press, 1990.
- [Wagner and Magyarik 85] N. R. Wagner and M. R. Magyarik. “A Public Key Cryptosystem Based on the Word Problem.” In *Advances in Cryptology. Proceedings of CRYPTO 1984*, pp. 19–36, edited by G. R. Blakley and D. Chaum, Lecture Notes in Computer Science 196. Berlin: Springer, 1985.

María Isabel González Vasco, Departamento de Matemáticas, Universidad de Oviedo, c/Calvo Sotelo, s/n, 33007 Oviedo, Spain (mvasco@orion.ciencias.uniovi.es)

Martin Rötteler, Centre for Applied Cryptographic Research, Department of Combinatorics and Optimization, University of Waterloo, Waterloo, Ontario, Canada, N2L 3G1 (mroetteler@cacr.math.uwaterloo.ca)

Rainer Steinwandt, Institut für Algorithmen und Kognitive Systeme, Arbeitsgruppe Computeralgebra, Prof. Dr. Th. Beth, Universität Karlsruhe, 76128 Karlsruhe, Germany (steinwan@ira.uka.de)

Received March 25, 2003; accepted April 17, 2003.